

Which components of heart failure programmes are effective? A systematic review and meta-analysis of the outcomes of structured telephone support or telemonitoring as the primary component of chronic heart failure management in 8323 patients: Abridged Cochrane Review[†]

Sally C. Inglis¹, Robyn A. Clark², Finlay A. McAlister³, Simon Stewart¹,
and John G.F. Cleland^{4*}

¹Preventative Health, Baker IDI Heart and Diabetes Institute and Faculty of Nursing, Midwifery and Health, University of Technology, Sydney, Australia; ²School of Nursing and Midwifery, Queensland University of Technology, Queensland, Australia; ³Division of General Internal Medicine, University of Alberta, Edmonton, Canada; and ⁴Academic Unit of Cardiology, Castle Hill Hospital, East Yorkshire, UK

Received 28 October 2010; accepted 9 November 2010; online publish-ahead-of-print 6 July 2011

Aims

Telemonitoring (TM) and structured telephone support (STS) have the potential to deliver specialized management to more patients with chronic heart failure (CHF), but their efficacy is still to be proven. The aim of this meta-analysis was to review randomized controlled trials (RCTs) of TM or STS for all-cause mortality and all-cause and CHF-related hospitalizations in patients with CHF, as a non-invasive remote model of a specialized disease-management intervention.

Methods and results

We searched all relevant electronic databases and search engines, hand-searched bibliographies of relevant studies, systematic reviews, and meeting abstracts. Two reviewers independently extracted all data. Randomized controlled trials comparing TM or STS to usual care in patients with CHF were included. Studies that included intensified management with additional home or clinic-visits were excluded. Primary outcomes (mortality and hospitalizations) were analysed; secondary outcomes (cost, length of stay, and quality of life) were tabulated. Thirty RCTs of STS and TM were identified (25 peer-reviewed publications ($n = 8323$) and five abstracts ($n = 1482$)). Of the 25 peer-reviewed studies, 11 evaluated TM (2710 participants), 16 evaluated STS (5613 participants) with two testing both STS and TM in separate intervention arms compared with usual care. Telemonitoring reduced all-cause mortality {risk ratio (RR) 0.66 [95% confidence interval (CI) 0.54–0.81], $P < 0.0001$ } and STS showed a similar, but non-significant trend [RR 0.88 (95% CI 0.76–1.01), $P = 0.08$]. Both TM [RR 0.79 (95% CI 0.67–0.94), $P = 0.008$], and STS [RR 0.77 (95% CI 0.68–0.87), $P < 0.0001$] reduced CHF-related hospitalizations. Both interventions improved quality of life, reduced costs, and were acceptable to patients. Improvements in prescribing, patient-knowledge and self-care, and functional class were observed.

Conclusion

Telemonitoring and STS both appear effective interventions to improve outcomes in patients with CHF. Systematic Review Number: Cochrane Database of Systematic Reviews. 2008:Issue 3. Art. No.: CD007228. DOI: 10.1002/14651858.CD007228.

[†] This paper is based on a Cochrane review first published in The Cochrane Database of Systematic Reviews (CDSR) 2010, Issue 8 (see www.thecochranelibrary.com for information). Cochrane reviews are regularly updated as new evidence emerges and in response to feedback, and the CDSR should be consulted for the most recent version of the review.

* Corresponding author. Tel: +44 1482 461776, Fax: +44 1482 461779, Email: j.g.cleland@hull.ac.uk

Published on behalf of the European Society of Cardiology. All rights reserved. © 2011 Cochrane collaboration, reproduced with permission. For permissions please email: journals.permissions@oup.com.

Keywords

Systematic review • Meta-analysis • Telemonitoring • Heart failure

Introduction

Chronic heart failure (CHF) specialized disease management programmes improve survival and quality of life, reduce rehospitalization, and increase the implementation of evidence-based practice.^{1,2} However, most of the successful CHF disease management programmes have been built around close clinical follow-up. The need for intense face-to-face follow-up strategies limits the number of patients who can participate in these programmes.

Table 1 Inclusion and exclusion criteria.

Inclusion criteria	
Study type	Randomized controlled trials
Publication	Full peer-reviewed publication (primary meta-analysis of primary outcomes)
Participants	Participants with a definitive diagnosis of heart failure, aged ≥ 18 years Recently discharged from an acute care setting to home (excluding nursing homes or convalescent homes) or recruited while managed in the community setting
Intervention	Structured scheduled telephone support or telemonitoring (daily, weekly, and monthly) Initiated by a healthcare professional (medical, nursing, social work, pharmacist). Delivered as the only heart failure disease management intervention, without home-visits or intensified clinic follow-up Targeted towards the patient, and not caregivers Did not include any visits at home by a specialized CHF healthcare professional or study personnel for the purpose of education or clinical assessment other than an initial visit to set-up equipment
Comparison	Consisted of standard post-discharge care without intensified attendance at cardiology clinics or clinic-based CHF disease management programme or home-visits
Outcomes	All-cause mortality, CHF-related or all-cause hospitalizations, length of stay, cost of the intervention or cost reductions, quality of life, acceptability, and adherence
Exclusion criteria	
No primary or secondary outcomes of interest reported or available from the study authors	
Not specific to heart failure	
Studies could not include any home-visits by specialized CHF health professionals or study personnel for the purpose of education or clinical assessment or include intensified clinic follow-up	

Copyright Cochrane Collaboration, reproduced with permission.

Structured telephone support (STS) is monitoring and/or self-care management delivered using simple telephone technology (data may have been collected and stored by a computer). Telemonitoring (TM) is digital/broadband/satellite/wireless, or blue-tooth transmission of physiological data e.g. electrocardiogram, blood pressure, weight, pulse oximetry, respiratory rate, and other data (self-care, education, lifestyle modification, and medicine administration). Both models of care have the potential to provide access to specialist care for a much larger number of patients across a much greater geography and might reduce the costs of care. These systems can assist directly in patient management, transferring the burden of care from health professionals and involve the patient in supported self-care.^{3,4} However, it is still not clear as to whether or not these interventions when delivered as the sole disease management intervention improve patient outcomes.^{5,6}

This review, published in the Cochrane Database of Systematic Reviews 2010⁷ updates a previously published review of remote monitoring strategies for CHF that included 10 trials of STS and five of TM.⁵ Since the first review, a number of large trials have been published reporting outcomes on both STS and TM. We have focused on the same primary outcomes (all-cause mortality, CHF-related hospitalization, and all-cause hospitalization) and secondary outcomes: length of stay, health-related quality of life, healthcare costs, and acceptability of the intervention to patients with CHF. Specifically, we have examined the benefits of STS or TM on a number of important outcomes in patients with CHF when compared with standard care, where STS or TM is the primary model of specialized disease-management intervention.

Methods

As per our protocol⁸ we applied the Cochrane methodology⁹ for this review.⁷ The specific eligibility criteria of included studies are presented in *Table 1*.

Information sources and search strategies

As per the Cochrane Heart Group protocol,⁹ all known relevant search engines and electronic databases were utilized for the review update period from January 2006 to November 2008. These included CENTRAL; MEDLINE; EMBASE; CINAHL; AMED; Science Citation Index Expanded; DARE; no date limit; National Research Register; IEEE Xplore; OALster; Google Scholar; Informit; Vivisimo; Australian Digital Theses Programme, and Proquest Digital Dissertations. Bibliographies of relevant studies and systematic reviews were hand-searched. Abstracts from the following conferences were also hand-searched for the years 2006, 2007, and 2008: European Society of Cardiology (ESC) Congress; American College of Cardiology Annual Scientific Sessions; American Heart Association Scientific Sessions; Heart Failure Society of America Annual Scientific Meeting; ESC Heart Failure Congress; ESC Spring Meeting of Cardiovascular Nursing; World Congress of Cardiology (2006, 2008); and the Asia-Pacific Heart Failure Congress (2008).

Keywords included: heart failure, cardiac failure, telehealth, telephone, telemonitoring and disease-management. Language restrictions were not applied. Full details of search strategies are available.⁷

Study selection

Randomized controlled trials (RCT) of STS or TM compared with usual care were eligible to be included in the meta-analysis if they were published in full in a peer-reviewed journal.⁹ Studies that were published as abstracts only were included in sensitivity meta-analyses.⁹ This decision was supported by a publication, co-authored by a member of this review team, which demonstrated that substantial potential discrepancies between results presented in meeting abstracts compared with final peer-reviewed publications.¹⁰

Data collection process

Two expert reviewers (S.C.I. and R.A.C.) independently reviewed the results of each search according to the inclusion and exclusion criteria with a standardized data extraction tool and also applied standard scales to judge study quality and risk of bias.⁹ A third reviewer (J.G.F.C.) adjudicated.⁹

Data and analysis

Meta-analyses of the primary outcomes (all-cause mortality, CHF-related, and all-cause hospitalizations) were performed according to Mantel-Haenszel methods, using a fixed effects model, risk ratios (RR), intention-to-treat, and assessment of statistical heterogeneity using the I^2 statistic.⁹ All analyses were performed using Review Manager (RevMan) Version 5.0 (Copenhagen: The Nordic Cochrane Centre, Cochrane Collaboration, 2008).⁹ Secondary outcomes such as quality of life, cost effectiveness and adherence and acceptance were measured using multiple tools across the studies. To summarize these outcomes results have been tabulated and described. Due to variances in the way length of stay was calculated and reported in the included studies, this outcome was tabulated as opposed to pooled into a meta-analysis.

Sensitivity analysis

Data from included studies published only as abstracts were added to the meta-analyses of the primary outcomes to assess whether publication status made any difference to the result, including the level of heterogeneity. A second sensitivity analysis was performed to assess the impact of length of follow-up on the primary outcomes for full peer-reviewed publications only, excluding studies with a follow-up period of 6 months or less.

Results

Study selection

Overall, 322 publications from 7952 citations were identified as potentially relevant studies and full copies were retrieved and assessed. Exclusions are detailed in *Figure 1*.

Study characteristics

Thirty RCTs of STS and TM were identified, which included 9560 participants (*Table 2*). These include 16 studies of STS^{14–18,21–23,25–30,32,33} (two of which were published abstracts^{14,23}) 12 studies of TM^{34–36,38–45,47} (three of which were published abstracts^{36,43,47}), and two studies with both

STS and TM intervention arms compared with usual care.^{11,13} The included populations were from seven countries (*Table 2*).

Twenty-five studies were published as full peer-reviewed publications (16 STS, $n = 5613$ and 11 TM, $n = 2710$). Two studies^{11,13} had two separate intervention arms (STS vs. TM vs. usual care) and each was considered as a separate comparison with usual care (and are included in the aforementioned counts). One study included two STS arms, one of which used standard telephone equipment and the other a videophone; for our analyses these two intervention arms were combined as STS.³³

Risk of bias

Analysis of the distribution in the funnel plots (not shown) demonstrated a strong publication bias towards positive outcomes in the included studies.^{7,9} A summary of the risk of bias analysis is presented in *Table 3*. The heterogeneity within the studies ranged from low (for all-cause mortality, $I^2 = 0\%$) to substantial (for all-cause hospitalizations, STS $I^2 = 24\%$; TM $I^2 = 78\%$) (*Figures 2–4*) (I^2 statistic low = 0–40%; moderate = 30–60%; substantial = 50–90%, considerable = 75–100%).⁹

All-cause mortality

Fifteen studies of STS^{11,13,15–18,21,22,25,26,28–30,32,33} and 11 studies of TM^{11,13,34,35,38–42,44,45} measured the effect on the risk of all-cause mortality (*Figure 2*). Telemonitoring was effective in reducing the risk of all-cause mortality in patients with CHF, with a 34% reduction in the risk of mortality observed [RR 0.66 95% confidence interval (CI) 0.54–0.81, $P < 0.0001$, $I^2 = 0\%$]. A similar non-significant trend was noted for STS [RR 0.88 (95% CI 0.76–1.01), $P = 0.08$, $I^2 = 0\%$].

Addition of studies published as abstracts^{14,23,36,43,47} had no substantial impact on the results [STS: RR 0.85, (95% CI 0.75–0.97), $I^2 = 0\%$, $P = 0.02$; TM: RR 0.68, (95% CI 0.57–0.82), $I^2 = 0\%$, $P < 0.0001$] nor did exclusion of studies lasting 6 months or less^{15,21,25,28,29,32,41,44,45} [STS: RR 0.87, (95% CI 0.74–1.02), $I^2 = 0\%$, $P = 0.08$; TM: RR 0.69, (95% CI 0.55–0.86), $P = 0.0009$, $I^2 = 0\%$].

Chronic heart failure-related hospitalization

Thirteen studies of STS^{11,13,15,16,21,22,25–30,32} and four studies of TM^{11,13,40,42} examined the effect of these interventions on the risk of CHF-related hospitalization (*Figure 3*). Structured telephone support reduced the proportion of patients hospitalized due to CHF by 23% [RR 0.77, (95% CI 0.68–0.87), $P < 0.0001$, $I^2 = 7\%$] and TM reduced this by 21% [RR 0.79, (95% CI 0.67–0.94), $P = 0.008$, $I^2 = 39\%$].

Addition of one STS study published as an abstract²³ did not alter the result of the meta-analysis other than to reduce heterogeneity ($I^2 = 0\%$). Addition of two TM studies published as abstracts^{43,47} to the meta-analysis marginally improved the effect [RR 0.76, (95% CI 0.64–0.89), $P = 0.0006$; $I^2 = 34\%$]. Excluding studies with a follow-up period of 6 months or less^{15,21,25,27–29,32} did not change the results of the meta-analyses [STS: RR 0.76, (95% CI 0.65–0.89), $P = 0.0005$; $I^2 = 0\%$]. All TM studies reporting CHF-related hospitalizations had a follow-up period > 6 months.




Figure 1 Study selection flowchart.⁷ Copyright Cochrane Collaboration, reproduced with permission.

All-cause hospitalization

Eleven studies of STS^{11,13,16,21,22,25,28–30,32,33} and eight studies of TM^{11,13,34,40–42,44,45} examined the effect of these interventions on the proportion of participants with at least one hospitalization for any cause (Figure 4). The effect of STS and TM on the risk of a hospitalization was similar [STS: RR 0.92, (95% CI 0.85–0.99), $P = 0.02$, $I^2 = 24\%$; TM: RR 0.91, (95% CI 0.84–0.99), $P = 0.02$, $I^2 = 78\%$].

No substantial change was observed in the results of the meta-analysis when studies published as abstracts were added^{23,43,47} [STS: RR 0.90 (95% CI 0.84–0.97), $P = 0.003$, $I^2 = 32\%$; TM: RR 0.94, (95% CI 0.87–1.01), $P = 0.09$, $I^2 = 73\%$]. Excluding studies with a follow-up period of 6 months or less,^{21,25,28,29,32,41,44} did not substantially change the outcome [STS: RR 0.91, (95% CI 0.83–0.99), $P = 0.03$; $I^2 = 22\%$; TM: RR 0.87, (95% CI 0.80–0.95), $P = 0.002$, $I^2 = 85\%$].

Table 2 Description of included studies.

Study	Intervention	Country	Duration of follow-up	Number of patients	Mean age (year)	Male sex (%)	Outcomes
Structured telephone support vs. usual care; telemonitoring vs. usual care							
Cleland et al. (2005) TEN-HMS study ^{11,12}	STS: education and monitoring.	Germany, Netherlands, UK	240 days; 15 months	426	67	77	Mortality, hospitalizations, length of stay, adaptation, and acceptance of the intervention
Mortara et al. (2009) HHH study ¹³	TM: weight, BP, ECG STS: education and monitoring TM: weight, BP and symptoms	UK, Poland, Italy	11.6 months (mean)	461	60	85	Mortality, hospitalizations, bed-days, adherence to the intervention
Structured telephone support vs. usual care							
Angermann et al. (2007) INH study [Abstract] ¹⁴	Education and monitoring	Germany	6 months	708	68	71	Mortality, days alive and out of hospital, NYHA functional class, and quality of life
Barth (2001) ¹⁵	Education and monitoring	USA	3 months	34	75	47	Mortality, unexpected visits, hospitalizations, cost, and quality of life
DeBusk et al. (2004) ¹⁶	CHF lifestyle education and medication management	USA	12 months	462	72	51	Mortality, hospitalizations, emergency and outpatient visits, prescription of recommended pharmacotherapy
DeWalt et al. (2006) ¹⁷	Education and monitoring	USA	12 months	127	62.5	47	Mortality, hospitalizations, quality of life, heart failure self-efficacy, heart failure knowledge, weight monitoring
Galbreath et al. (2004) ¹⁸⁻²⁰	Education and monitoring	USA	18 months	1069	71	71	Mortality, 6-min walk test, NYHA functional class, quality of life, and cost. Subgroup: ejection fraction and medication adherence
Gattis et al. (1999) PHARM study ²¹	Pharmacist-led medication review and education	USA	6 months	181	67	68	Mortality, hospitalization and medication prescription
GESICA Investigators (2005) DIAL trial ²²	Education and monitoring	Argentina	16 months (mean)	1518	65	71	Mortality, hospitalizations, quality of life, and medication prescription
Tonkin et al. (2009) CHAT study ^{23,24} [Abstract]	Telewatch system (Baltimore)	Australia	12 months	405	NR	NR	Mortality, hospitalizations, adherence to, adaptation, and acceptance of intervention
Laramee et al. (2003) ²⁵	Education and monitoring	USA	3 months	287	71	54	Mortality, hospitalizations, costs, medication prescription, and adherence
Rainville (1999) ²⁶	Pharmacist-led medication review, education, medication management	USA	12 months	38	70	50	Mortality, hospitalizations, and NYHA functional class
Ramachandran et al. (2007) ²⁷	Education, monitoring, and medication management	India	6 months	50	44.5	78	Hospitalizations, NYHA functional class, quality of life, medication prescription, and cost
Riegel et al. (2002) ²⁸	Education and counselling	USA	6 months	358	74	49	Mortality, hospitalizations, hospital days, physician and emergency department visits, cost, and patient satisfaction
Riegel et al. (2006) ²⁹	Education, monitoring, and guidance	USA	6 months	135	72	46	Mortality, hospitalizations, cost, quality of life, and depression
Sisk et al. (2006) ^{30,31}	Patient assessment and education	USA	12 months	406	59	54	Mortality, hospitalizations, cost, and quality of life

Tsuyuki <i>et al.</i> (2004) ³²	Education and monitoring	Canada	6 months	276	72	58	Mortality, hospitalizations, medication adherence, physician and emergency department visits, and cost
Wakefield <i>et al.</i> (2008) ³³	Education and monitoring	USA	12 months	148	69	99	Mortality, hospitalizations, hospital days, time to first readmission, urgent care clinic visits, quality of life, satisfaction, and cost
Telemonitoring vs. usual care							
Antoniceilli <i>et al.</i> (2008) ³⁴	BP, HR, weight and 24h urine output, ECG	Italy	12 months	57	78	61	Mortality, hospitalizations, and quality of life
Balk <i>et al.</i> (2008) ³⁵	MOTIVA system	The Netherlands	288 days (mean)	214	66	70	Hospital days, days alive and out of hospital, quality of life, heart failure knowledge, cost, acceptance, and self-care
Blum <i>et al.</i> (2007) [Abstract] ^{36,37}	Weight, BP, HR, rhythm	USA	12 months	204	72	71	Hospitalizations, quality of life, mortality, and BNP
Capomolla <i>et al.</i> (2004) ³⁸	Weight, systolic BP, HR	Italy	12 months	133	57	88	Mortality, hospitalizations, adherence, and emergency department visits
de Lusignan <i>et al.</i> (2001) ³⁹	Pulse, BP, weight	UK	12 months	20	75	NR	Mortality, satisfaction, adherence, and quality of life
Giordano <i>et al.</i> (2009) ⁴⁰	One-lead ECG	Italy	12 months	460	57	85	Mortality, hospitalizations, haemodynamic instability episode occurrence, and cost
Goldberg <i>et al.</i> (2003) WHARF trial ⁴¹	Weight and symptoms	USA	6 months (mean)	280	59	68	Mortality, hospitalizations, emergency department visits, quality of life, and adherence to the intervention
Kielblock <i>et al.</i> (2007) ⁴²	Weight	Germany	12 months	502	74	51	Mortality, length of stay, hospital and drug costs, total costs per patient, satisfaction, hospitalizations, and medication prescription
Villani <i>et al.</i> (2007) ICARUS [Abstract] ⁴³	Weight, urine output, fluid intake, BP, HR	Italy	12 months	77	69	75	Mortality, hospitalizations, emergency room visits, and hospital days per patient
Soran <i>et al.</i> (2008) ⁴⁴	Weight and symptoms	USA	6 months	315	76	35	Mortality, hospitalizations, length of hospital stay, emergency room visits, quality of life, and adherence to the intervention
Woodend <i>et al.</i> (2008) ^{45,46}	Weight, ECG, BP	Canada	3 and 12 months	121	68	74	Mortality, hospitalizations, quality of life, emergency department visits, hospital days, and patient satisfaction
Zugck <i>et al.</i> (2008) HiTel trial [Abstract] ⁴⁷	Weight, BP, 12-lead ECG	Germany	12 months	88	58	82	Mortality, hospitalizations, and length of stay

Table 3 Assessment of bias of included studies.

Study	Adequate sequence generation	Adequate allocation concealment	Adequate blinding of outcome assessors	Incomplete outcome data assessed	Free of selective outcome reporting
Structured telephone support vs. usual care; telemonitoring vs. usual care					
Cleland et al. (2005) TEN-HMS study ¹¹	Yes	Yes	Unclear	Yes	Yes
Mortara et al. (2009) HHH study ¹³	Unclear	Yes	Yes	Unclear	Unclear
Structured telephone support vs. usual care					
Angermann et al. (2007) INH study [Abstract] ¹⁴	Unclear	Unclear	Unclear	Unclear	Unclear
Barth (2001) ¹⁵	Unclear	Unclear	Unclear	Unclear	Unclear
DeBusk et al. (2004) ¹⁶	Yes	Yes	Yes	Yes	Yes
DeWalt et al. (2006) ¹⁷	Yes	Unclear	No	Yes	Yes
Galbreath et al. (2004) ¹⁸	Unclear	Yes	No	No	Yes
Gattis et al. (1999) PHARM study ²¹	Yes	No	Yes	Unclear	Yes
GESICA Investigators (2005) DIAL trial ²²	Yes	Yes	Yes	Yes	Yes
Tonkin et al. (2009) CHAT study ²³ [Abstract]	Unclear	Unclear	Unclear	Unclear	Unclear
Laramée et al. (2003) ²⁵	Unclear	Unclear	Unclear	Unclear	No
Rainville (1999) ²⁶	Unclear	Yes	Unclear	Yes	Yes
Ramachandran et al. (2007) ²⁷	Yes	Unclear	Unclear	Unclear	Unclear
Riegel et al. (2002) ²⁸	Unclear	Yes	Unclear	Unclear	Yes
Riegel et al. (2006) ²⁹	Unclear	Yes	Yes	Yes	Yes
Sisk et al. (2006) ³⁰	Yes	Yes	Unclear	Yes	Yes
Tsuyuki et al. (2004) ³²	Unclear	Unclear	Unclear	Unclear	Yes
Wakefield et al. (2008) ³³	Yes	Yes	Unclear	Yes	Yes
Telemonitoring vs. usual care					
Antonicelli et al. (2008) ³⁴	Unclear	Unclear	Unclear	Yes	Yes
Balk et al. (2008) ³⁵	Yes	Unclear	Unclear	Yes	Yes
Blum et al. (2007) [Abstract] ³⁶	Yes	Yes	Unclear	Yes	Unclear
Capomolla et al. (2004) ³⁸	Unclear	Unclear	Unclear	Unclear	Yes
de Lusignan et al. (2001) ³⁹	Yes	Unclear	Unclear	Unclear	Yes
Giordano et al. (2009) ⁴⁰	Unclear	Unclear	Unclear	Yes	Yes
Goldberg et al. (2003) WHARF trial ⁴¹	Unclear	Yes	Yes	Yes	Unclear
Kielblock et al. (2007) ⁴²	No	Unclear	Unclear	Unclear	Unclear
Villani et al (2007) ICARUS [Abstract] ⁴³	Yes	Unclear	Unclear	Unclear	Unclear
Soran et al. (2008) ⁴⁴	Unclear	Unclear	Yes	Unclear	Yes
Woodend et al. (2008) ⁴⁵	Unclear	Unclear	Unclear	Unclear	No
Zugck et al. (2008) HiTel trial [Abstract] ⁴⁷	Unclear	No	Unclear	Unclear	Unclear

Copyright Cochrane Collaboration, reproduced with permission.


Length of stay

Only one STS study³² reported a statistically significant reduction in the length of stay for patients in the intervention group compared with those receiving usual care. One study reported a substantial difference in the number of hospital days per patient.⁴³

Health-related quality of life

Sixteen studies reported quality of life.^{14,15,17,19,22,27,29,30,33–35,37,39,41,44,45} These were either a direct comparison between the intervention and control groups at study conclusion, or between baseline and study conclusion within the study arm. A range of psychometric

Structured telephone support



Telemonitoring




Figure 2 Effect of structured telephone support and telemonitoring on all-cause mortality.⁷ Copyright Cochrane Collaboration, reproduced with permission.

tools were used [Chronic Heart Failure Symptomatology Questionnaire; Minnesota Living with Heart Failure Questionnaire (MLWHFQ); Kansas City Cardiomyopathy Questionnaire (KCCM); Short-Form-12 Item; Short-Form-36 Item (SF-36); Health Distress Score].

Six studies of STS^{14,15,22,27,30,33} reported improvements in quality of life, with significant improvements in physical ($P = 0.03$)¹⁴ and overall measures (MLWHFQ and KCCM). Three TM studies,^{34,37,45,46} reported improvements in quality of life (MLWHFQ $P = 0.001$ and SF-36 mental ($P = 0.001$), and physical component scores ($P = 0.003$),³⁷ MLWHFQ $P = 0.025$,⁴⁶ and SF-36 $P < 0.05$,⁴⁵ SF-36 health perception $P = 0.046$ ³⁴).

Cost


Twelve studies (nine STS^{15,18,20,25,27–29,31–33} and three TM^{35,40,42}) provided details on cost of the intervention or cost reductions associated with the intervention or cost effectiveness. The cost

of the interventions varied according to the type of intervention, in particular the technologies used and the intensity at which it was delivered. Of the 11 studies which reported the effect of the intervention on the cost of care,^{18,20,25,27–29,32,33,35,40,42} all but three,^{18,20,29,35} reported reductions in cost (either cost per admission or overall reduction in healthcare costs), with those reporting per cent reductions ranging between 14%²⁵ and 86%.³³

Other outcomes

Few studies reported adherence to the intervention (compliance). Among those that did,^{13,24,38,39,41,44} adherence was measured at 65.8% for STS,²⁴ and 75% to 98.5% for TM.^{13,38,39,41,44} The adaptation to the technology was high, with two studies, reporting that 96–97% of patients (often aged >70 years) were able to learn and use the STS or TM systems.^{12,24} Acceptance (satisfaction) of patients receiving healthcare via STS or TM was rated

Structured telephone support



Telemonitoring




Figure 3 Effect of structured telephone support and telemonitoring on CHF-related hospitalizations.⁷ Copyright Cochrane Collaboration, reproduced with permission.

between 76%²⁴ and 100%.^{12,24,35,42,45} Improvements in other outcomes from these trials included: New York Heart Association (NYHA) functional class that improved in three^{14,18,27} studies. Chronic heart failure knowledge and self-care improved in both studies reporting this outcome;^{17,35} 6 min walk test improved in one study,²⁷ of the two¹⁸ that reported this outcome; improvements in evidence-based pharmacotherapy were seen in six^{18,21,22,25,27,42} of the seven studies¹⁶ that reported this outcome and the only study to report brain natriuretic peptide, reported an improvement in this measure.³⁷

Discussion

This systematic review and meta-analysis suggests that both TM and STS have a broad range of benefits for patients with heart failure: including a substantial reduction in all-cause mortality for TM, a substantial reduction in the risk of CHF hospitalization for both TM and STS and a modest reduction in the risk of all-cause hospitalization. These interventions improved quality of life, reduced costs, and were acceptable to patients. Improvements in prescribing, patient knowledge and self-care, and functional class were observed. The precise mechanisms by which these

interventions produce these effects are unclear but probably reflect a combination of improved implementation of and adherence to guideline therapies, early identification of complications or disease progression, and a positive impact on patient psychology.⁴⁸ Patients in these trials reported a sense of reassurance and security, feeling that they have a lifeline to expert care.^{23,27}

Two substantial studies^{49,50} have been reported since this review was completed and will be incorporated in the next revision which is currently underway. The Tele-HF study⁴⁹ which included 1653 patients, was a study of a voice interactive system (STS) applied to patients recently discharged from hospital after an episode of worsening heart failure. Adherence with the system was very poor, suggesting that patients did not engage with the service, perhaps because of the nature of the technology. No benefits were observed on death or hospitalization. This is consistent with the results of our systematic review,^{7,51} at least with respect to mortality. The TIM-HF study,^{50,52,53} including 710 patients, was a TM study of patients with exceptionally well-managed chronic stable heart failure monitored by a remote expert group. Trends to fewer deaths and hospitalizations with TM were not significant; suggesting that home TM might not be an effective intervention in stable patients when other systems



Structured telephone support**Telemonitoring**

Figure 4 Effect of structured telephone support and telemonitoring on all-cause hospitalizations.⁷ Copyright Cochrane Collaboration, reproduced with permission.

have ensured a high quality of care. However, TM might be a more efficient and less expensive option when the quality of care is not of a similar standard to that provided in TIM-HF. Also, it is unclear as to whether home TM is most successfully deployed as an adjunct to personalized care from a local specialist clinic or as a remote regional or national service.

We excluded from this analysis other methods of follow-up and management that have also been reported to improve outcomes, such as nurse-led or specialist heart failure clinic or home visits.²

It is possible that intensified self-management and remote management are the key factors driving clinical benefit with these interventions. If so, the main issues revolve around an organization and cost effectiveness. The most expensive aspect of healthcare in high-income countries is staff to run services and deliver care. Delivering care by increasing direct one-to-one interactions is likely to be an expensive long-term strategy. Development of TM systems that support the patient directly in making decisions about issues such as diuretic dose, diet, and life-style and when to seek professional advice have the potential to offer expert care to most patients with CHF. Implementation of TM will require a change in the approach of healthcare systems to the delivery of care

with redeployment of existing staff rather than an expansion of the healthcare work force required by other strategies. Indeed, it is quite likely that TM has not worked optimally in clinical trials since the studies were generally done in parallel to rather than integrated with existing services. Restructuring healthcare around TM could be more effective and cost efficient.

An additional and increasingly apparent dimension to TM is that it is a direct investment in the patient rather than in healthcare services. The patient is less likely to be a passive recipient of services from health professionals, and becomes more actively involved in their care. Patients provide information on symptoms and vital signs and receive feedback and education, which they can review at their leisure as often as they wish and together with their carers and family. More advanced systems will ensure that the patients know when and how medication should be adjusted and when they can do this themselves and when they need professional support. Because the patients know what care they should receive, health professionals may be more likely to deliver it or explain why the patients should deviate from the plan. Telemonitoring will create more expert patients. Undoubtedly, this will create headaches for health professionals, leading to resistance to change.

The medical profession should offer the best service to patients even if this means moving out of their professional comfort zone.

Compared with other recent systematic reviews of remote monitoring in CHF,^{54,55} our review⁷ is unique in using robust Cochrane methodology.^{8,9} We have synthesized and quantified the benefits of STS and TM, while limiting the influence of confounders such as home-visits by specialized healthcare staff or frequent visits to a specialized CHF clinic on the efficacy of these interventions in managing patients with CHF. Previous reviews on this topic have included a mixture of research methods (RCTs and cohort studies) and studies of both invasive and non-invasive remote monitoring, many of which have involved home visits.^{54,55}

These findings have important clinical implications. The findings of this review are highly relevant to the future planning and implementation of CHF disease-management globally. This analysis provides strong evidence that these technologies reduce mortality and hospitalizations as well as improving measures such as quality of life. There may be benefits of using these technologies to manage patients with CHF that relate to human or financial resources, but perhaps the biggest advantage can be gained from utilizing these technologies to reach patients with CHF who are without access to home or clinic-based CHF-management programmes.^{3,4} Such benefits may not be restricted to high-income countries. Indeed, China and India⁵⁶ both have programmes for delivering care remotely.

The average age of patients in these included trials ranged from 45 to 78 years, with the majority of patients aged >68 years. It is clear that many older people are able to use and benefit from STS and TM. In fact TM devices are usually designed specifically with older people in mind.

Our synthesis of the evidence of STS and TM is only as good as the included studies. We were limited by the format of published results, especially for those where we were unsuccessful in obtaining further study details. In addition, the heterogeneity is large for some of the meta-analyses of the primary outcomes. This heterogeneity is not only within methodology but also the types and intensity of applied technologies.

There was evidence of publication bias. It is likely that many small studies are never published, either because the investigator does not offer their results for publication or because editors reject under-powered and negative studies. There is a dearth of evidence about how long patients should be supported by TM or STS. It is possible that the greatest benefit in terms of education and medication patterns is accrued within a few weeks and that long-term monitoring is redundant. However, the weight monitoring in heart failure trial showed that a 6-month TM intervention was associated with a reduction in mortality but that withdrawal led to rapid loss of this initial benefit, suggesting that long-term TM might be superior to short-term TM.⁵⁷

In conclusion, STS and TM improve outcomes for patients with CHF, although only TM appears to have a substantial impact on reducing mortality. This may reflect the impact of improved access to specialist care, which could be delivered by more conventional means but at additional cost. The effects appear substantial and might be an underestimate of the true impact when properly integrated into care pathways. Given the wealth of

evidence and the impact on hospitalization that is likely to mitigate costs, all patients with CHF should have access to enhanced surveillance.

Acknowledgements

This paper is based on a Cochrane review first published in The Cochrane Database of Systematic Reviews (CDSR) 2010, Issue 8 (see www.thecochranelibrary.com for information). Cochrane reviews are regularly updated as new evidence emerges and in response to feedback, and the CDSR should be consulted for the most recent version of the review.

We wish to acknowledge the valuable contribution of the following researchers who contributed to this review: Dr Christian Lewinter and Dr Damien Cullington from academic unit of cardiology, Castle Hill Hospital, East Yorkshire, United Kingdom for retrieving studies and undertaking hand-searching; Ms Jocasta Ball from Baker IDI Heart and Diabetes Institute, Melbourne, Australia for assistance with retrieving studies and entering study details into a bibliography and Ms Stefanie Nagendirajah from Baker IDI Heart and Diabetes Institute, Melbourne, Australia for assistance with retrieving studies. We would also like to thank the staff at the Cochrane Collaboration Heart Review Group, in particular, Dr Joey Kwong, Managing Editor, and Miss Claire Williams, Assistant Managing Editor. We would like to also thank Ms Monika Winterstein, Mr Horst Winterstein, Mrs Erika Winterstein, and Ms Andrea Horsky for their assistance with German translations. Our team also wishes to make special acknowledgement to the contribution of the following librarians for their assistance with designing and conducting the searches: Ms Margaret Burke (Cochrane Heart Review Group Trial Search Coordinator), Dr Helen Marlborough (Medical Science Librarian, University of Glasgow); Ms Margaret Goedhart (Health Sciences Librarian, University of South Australia). We also acknowledge the following study authors who were very generous in sharing further details of their studies and data, some of which is unpublished, in order to include the most up-to-date data in our meta-analysis: CE Angermann, K Blum; DA DeWalt; M Blasius; P Brocki; S Kottmair; WA Gattis; LR Goldberg; A Laramee; A Mortarra; G Parati; B Riegel; RT Tsuyuki; BJ Wakefield; R Cebola and D Schellberg.

Funding

This study received no external funding. All authors had full access to all the data in the study and had final responsibility for the decision to submit for publication. S.C.I. is a Post-Doctoral Research Fellow supported by the National Health and Medical Research Council of Australia (NHMRC) and National Heart Foundation of Australia (NHMRC Grant ID 472 699). R.A.C. is a Post-Doctoral Research Fellow supported by the NHMRC (NHMRC Grant ID 570 141) and a Research SA Fellowship. S.S. is a Senior Research Fellow supported by the NHMRC (NHMRC Grant ID 472 658). F.A.M. receives salary support from the Alberta Heritage Foundation for Medical Research Health Scholar Program and the University of Alberta/Merck Frost/Aventis Chair in Patient Health Management.

Conflict of interest

J.G.F.C. has received funds from Philips and Bosch, which have a commercial interest in telemonitoring, for research staff and fees

for consulting and has acted as a paid advisor on the subject of this review. J.G.F.C. was involved in the design, conduct and publication of a study included in this review. S.S. was involved in the design and publication of a study included in this review. R.A.C. was involved in the conduct and publication of a study included in this review. No other funding was received to support this project.

References

- Gonseth J, Guallar-Castillon P, Banegas JR, Rodriguez-Artalejo F. The effectiveness of disease management programmes in reducing hospital re-admission in older patients with heart failure: a systematic review and meta-analysis of published reports. *Eur Heart J* 2004;**25**:1570–1595.
- McAlister FA, Stewart S, Ferrua S, McMurray J. Multidisciplinary strategies for the management of heart failure patients at high risk for admission—a systematic review of randomized trials. *J Am Coll Cardiol* 2004;**44**:810–819.
- Jaarsma T, Stromberg A, De Geest S, Fridlund B, Heikkilä J, Mårtensson J, Moons P, Scholte op Reimer W, Smith K, Stewart S, Thompson DR. Heart failure management programmes in Europe. *Eur J Cardiovasc Nurs* 2006;**5**: 197–205.
- Clark RA, Driscoll A, Nottage J, McLennan S, Coombe DM, Bamford EJ, Wilkinson D, Stewart S. Inequitable provision of optimal services for patients with chronic heart failure: a national geo-mapping study. *Med J Aust* 2007;**186**: 169–174.
- Clark RA, Inglis SC, McAlister FA, Cleland JGF, Stewart S. Telemonitoring or structured telephone support programmes for patients with chronic heart failure: systematic review and meta-analysis. *BMJ* 2007;**334**:942–945.
- Taylor S, Bestall J, Cotter S, Falshaw M, Hood S, Parsons S, Wood L, Underwood M. Clinical service organisation for heart failure. *Cochrane Database Syst Rev* 2005, Issue 2. Art. No. CD002752.
- Inglis SC, Clark RA, McAlister FA, Ball J, Lewinter C, Cullington D, Stewart S, Cleland JG. Structured telephone support or telemonitoring programmes for patients with chronic heart failure. *Cochrane Database Syst Rev* 2010, Issue 8. Art. No. CD007228.
- Inglis SC, Clark RA, Cleland JGF, McAlister F, Stewart S. Structured telephone support or telemonitoring programs for patients with chronic heart failure. *Cochrane Database Syst Rev* 2008, Issue 3. Art. No. CD007228.
- Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions Version 5.0.1 [updated September 2008]*. www.cochrane-handbook.org The Cochrane Collaboration, 2008.
- Toma M, McAlister FA, Bialy L, Adams D, Vandermeer B, Armstrong PW. Transition from meeting abstract to full-length journal article for randomized controlled trial. *JAMA* 2006;**295**:1281–1287.
- Cleland JG, Louis AA, Rigby AS, Janssens U, Balk AH. Non-invasive home telemonitoring for patients with heart failure at high risk of recurrent admission and death: the Trans-European Network-Home-Care Management System (TEN-HMS) study. *J Am Coll Cardiol* 2005;**45**:1654–1664.
- Louis AA, Balk A, Janssens U, Westerteicher C, Cleland JG. Patient acceptance and satisfaction of home telemonitoring in the management of heart failure: TENS-HMS (abstract). *JACC* 2002;**19**:S37A.
- Mortara A, Pinna GD, Johnson P, Maestri R, Capomolla S, La Rovere MT, Ponikowski P, Tavazzi L, Sleight P, HHH Investigators. Home telemonitoring in heart failure patients: the HHH study (Home or Hospital in Heart Failure). *Eur J Heart Fail* 2009;**11**:312–318.
- Angermann CE, Stork S, Gelbrich G, Faller H, Jahns R, Frantz S, Ertl G. A prospective randomized controlled trial comparing the efficacy of a standardized, supraclassically transferable program for monitoring and education of patients with systolic heart failure with usual care—the Interdisciplinary Network for Heart Failure (INH) Study (abstract). *Circ*, 2007;**116**(II):601.
- Barth V. A nurse-managed discharge program for congestive heart failure patients: outcomes and costs. *Home Health Care Manag Pract* 2001;**14**:436–43.
- DeBusk RF, Miller NH, Parker KM, Bandura A, Kraemer HC, Cher DJ, West JA, Fowler MB, Greenwald G. Care management for low-risk patients with heart failure: a randomized, controlled trial. *Ann Intern Med* 2004;**141**:606–613.
- DeWalt DA, Malone RM, Bryant ME, Kosnar MC, Corr KE, Rothman RL, Sueta CA, Pignone MP. A heart failure self-management program for patients of all literacy levels: a randomized, controlled trial. *BMC Health Serv Res* 2006;**6**:30.
- Galbreath AD, Krasuski RA, Smith B, Stajduhar KC, Kwan MD, Ellis R, Freeman GL. Long-term healthcare and cost outcomes of disease management in a large, randomized, community-based population with heart failure. *Circulation* 2004;**110**:3518–3526.
- Smith B, Forkner E, Zaslow B, Krasuski RA, Stajduhar K, Kwan M, Ellis R, Galbreath AD, Freeman GL. Disease management produces limited quality-of-life improvements in patients with congestive heart failure: evidence from a randomized trial in community-dwelling patients. *Am J Manag Care* 2005;**11**:701–713.
- Smith B, Hughes-Cromwick PF, Forkner E, Galbreath AD. Cost-effectiveness of telephonic disease management in heart failure. *Am J Manag Care* 2008;**14**: 106–115.
- Gattis WA, Hasselblad V, Whellan DJ, O'Connor CM. Reduction in heart failure events by the addition of a clinical pharmacist to the heart failure management team: results of the Pharmacist in Heart Failure Assessment Recommendation and Monitoring (PHARM) Study. *Arch Intern Med* 1999;**159**:1939–1945.
- GESICA Investigators. Randomised trial of telephone intervention in chronic heart failure: DIAL trial. *BMJ* 2005;**331**:425–427.
- Tonkin A, Yallop J, Driscoll A, Forbes A, Croucher J, Chan B, Stewart S, Clark R, Huynh L, Meehan A, Egan H, Piterman L, Kasper E, Krum H. Does telephone support of the rural and remote patients with heart failure improve clinical outcomes? Results of the Chronic Heart Failure Assistance by Telephone Study (CHAT) Study (abstract). *Heart Lung Circ* 2009;**18**:S105.
- Clark RA, Yallop JJ, Piterman L, Croucher J, Tonkin A, Stewart S, Krum H. CHAT Study Team. Adherence, adaptation and acceptance of elderly chronic heart failure patients to receiving healthcare via telephone-monitoring. *Eur J Heart Fail* 2007;**9**:1104–1111.
- Laramee AS, Levinsky SK, Sargent J, Ross R, Callas P. Case management in a heterogeneous congestive heart failure population: a randomized controlled trial. *Arch Intern Med* 2003;**163**:809–817.
- Rainville EC. Impact of pharmacist interventions on hospital readmissions for heart failure. *Am J Health Syst Pharm* 1999;**56**:1339–1342.
- Ramachandran K, Husain N, Maikhuri R, Seth S, Vij A, Kumar M, Srivastava N, Prabhakaran D, Airan B, Reddy KS. Impact of a comprehensive telephone-based disease management programme on quality-of-life in patients with heart failure. *Natl Med J India* 2007;**20**:67–73.
- Riegel B, Carlson B, Kopp Z, LePetri B, Glaser D, Unger A. Effect of a standardized nurse case-management telephone intervention on resource use in patients with chronic heart failure. *Arch Intern Med* 2002;**162**:705–712.
- Riegel B, Carlson B, Glaser D, Romero T. Randomized controlled trial of telephone case management in Hispanics of Mexican origin with heart failure. *J Card Fail* 2006;**12**:211–219.
- Sisk JE, Hebert PL, Horowitz CR, McLaughlin MA, Wang JJ, Chassin MR. Effects of nurse management on the quality of heart failure care in minority communities: a randomized trial. *Ann Intern Med* 2006;**145**:273–283.
- Hebert PL, Sisk JE, Wang JJ, Tuzzio L, Casabianca JM, Chassin MR, Horowitz C, McLaughlin MA. Cost-effectiveness of nurse-led disease management for heart failure in an ethnically diverse urban community. *Ann Intern Med* 2008;**149**: 540–548.
- Tsuyuki RT, Fradette M, Johnson JA, Bungard TJ, Eurich DT, Ashton T, Gordon W, Ikuta R, Kornder J, Mackay E, Manyari D, O'Reilly K, Semchuk W. A multicenter disease management program for hospitalized patients with heart failure. *J Card Fail* 2004;**10**:473–480.
- Wakefield BJ, Ward MM, Holman JE, Ray A, Scherubel M, Burns TL, Kienzle MG, Rosenthal GE. Evaluation of home telehealth following hospitalization for heart failure: a randomized trial. *Telemed J E Health* 2008;**14**:753–761.
- Antonicelli R, Testarmata P, Spazzafumo L, Gagliardi C, Bilo G, Valentini M, Olivieri F, Parati G. Impact of telemonitoring at home on the management of elderly patients with congestive heart failure. *J Telemed Telecare* 2008;**14**: 300–305.
- Balk AH, Davise W, Dommelen P, Klaassen E, Caliskan K, van der Burgh P, Leenders CM. Tele-guidance of chronic heart failure patients enhances knowledge about the disease. A multi-centre, randomised controlled study. *Eur J Heart Fail* 2008;**10**:1136–1142.
- Blum K, Gottlieb S. Morbidity and mortality benefits of reliable instrumental support (abstract). *J Cardiac Fail* 2007;**13**:S164.
- Blum K, Janowick F, Gottlieb SS. One year changes in quality of life for heart failure patients in a home telemonitoring program (abstract). *J Cardiac Fail*, 2006;**12**(Suppl 6):S122.
- Capomolla S, Pinna G, La Rovere MT, Maestri R, Ceresa M, Ferrari M, Febo O, Caporotondi A, Guazzotti G, Lenta F, Baldin S, Mortara A, Cobelli F. Heart failure case disease management program: a pilot study of home telemonitoring versus usual care. *Eur Heart J Suppl* 2004;**6**:F91–98.
- de Lusignan S, Wells S, Johnson P, Meredith K, Leatham E. Compliance and effectiveness of 1 year's home telemonitoring. The report of a pilot study of patients with chronic heart failure. *Eur J Heart Fail* 2001;**3**:723–730.
- Giordano A, Scalvini S, Zanelli E, Corra U, Longobardi GL, Ricci VA, Baiardi P, Glisenti F. Multicenter randomised trial on home-based telemanagement to prevent hospital readmission of patients with chronic heart failure. *Int J Cardiol* 2009;**131**:192–199.
- Goldberg LR, Piette JD, Walsh MN, Frank TA, Jaski BE, Smith AL, Rodriguez R, Mancini DM, Hopton LA, Orav EJ, Loh E, WHARF Investigators. Randomized

- trial of a daily electronic home monitoring system in patients with advanced heart failure: the Weight Monitoring in Heart Failure (WHARF) trial. *Am Heart J* 2003; **146**:705–712.
42. Kielblock B, Frye C, Kottmair S, Hudler T, Siegmund-Schultze E, Middeke M. Impact of telemetric management on overall treatment costs and mortality rate among patients with chronic heart failure. *Dtsch Med Wochenschr* 2007; **132**: 417–422.
 43. Villani A, Malfatto G, Della Rosa F, Branzi G, Boarin S, Borghi C, Cosentino E, Gualerzi M, Coruzzi P, Molinari E, Compare A, Cassi M, Collatina S, Parati G. Disease management for heart failure patients: role of wireless technologies for telemedicine. The ICAROS project (abstract). *G Ital Cardiol* 2007; **8**:107–114.
 44. Soran OZ, Pina IL, Lamas GA, Kelsey SF, Selzer F, Pilotte J, Lave JR, Feldman AM. A randomized clinical trial of the clinical effects of enhanced heart failure monitoring using a computer-based telephonic monitoring system in older minorities and women. *J Card Fail* 2008; **14**:711–717
 45. Woodend AK, Sherrard H, Fraser M, Stuewe L, Cheung T, Struthers C. Telehome monitoring in patients with cardiac disease who are at high risk of readmission. *Heart Lung* 2008; **37**:36–45.
 46. Woodend AK, Sherrard H, Fraser M, Stuewe L, Haddad H, Cheung T, Struthers C. Getting connected: telehome care for patients with heart disease. *Can Home Econ J* 2003; **52**:22–26.
 47. Zugck C, Frankenstein L, Nelles M, Froehlich H, Schellberg D, Cebola R, Remppis A, Katus HA. Telemedicine reduces hospitalisation rates in patients with chronic heart failure—results of the randomized HiTel trial (abstract). *Eur J Heart Fail Suppl* 2008; **7**:9a.
 48. Granger BB, Swedberg K, Ekman I, Granger CB, Olofsson B, McMurray JJ, Yusuf S, Michelson EL, Pfeffer MA, CHARM investigators. Adherence to candesartan and placebo and outcomes in chronic heart failure in the CHARM programme: double-blind, randomised, controlled clinical trial. *Lancet* 2005; **366**:2005–2011.
 49. Chaudhry SI, Mattera JA, Curtis JP, Spertus JA, Herrin J, Lin Z, Phillips CO, Hodshon BV, Cooper LS, Krumholz HM. Telemonitoring in patients with heart failure. *N Engl J Med* 2010; **363**:2301–2309.
 50. Kohler F, Winkler S, Schieber M, Sechtem U, Stangl K, Böhm M, Boll H, Gelbrich G, Kirwan B, Anker SD. Telemedical Interventional Monitoring in Heart Failure (TIM-HF), a Randomized, Controlled Intervention Trial Investigating the Impact of Telemedicine on Mortality in Ambulatory Patients With Chronic Heart Failure (abstract). *Circ* 2010; **122**:2215–2226.
 51. Inglis SC, Clark RA, Cleland JGF. Telemonitoring in patients with heart failure. Letter to the Editor. *N Engl J Med* 2011; **364**:1078–1079.
 52. Koehler F, Winkler S, Schieber M, Sechtem U, Stangl K, Bohm M, Boll H, Kim SS, Koehler K, Lucke S, Honold M, Heinze P, Schweizer T, Braecklein M, Kirwan B, Gelbrich G, Anker SD on behalf of the TIM-HF Investigators. Telemedical Interventional Monitoring in Heart Failure (TIM-HF), a randomized, controlled intervention trial investigating the impact of telemedicine on mortality in ambulatory patients with heart failure: study design. *Eur J Heart Fail* 2010; **12**: 1354–1362.
 53. Cleland JGF, Coletta AP, Buga L, Antony R, Pellicori P, Freemantle N, Clark AL. Clinical trials update from the American Heart Association Meeting 2010: EMPASIS-HF, RAFT, TIM-HF, Tele-HF, ASCEND-HF, ROCKET-AF and PROTECT. *Eur J Heart Fail* 2011 (April issue).
 54. Klersy C, De Silvestri A, Gabutti G, Regoli F, Auricchio A. Meta-Analysis of remote monitoring of heart failure patients. *J Am Coll Cardiol* 2009; **54**:1683–1694.
 55. Dang S, Dimmick S, Kelkar G. Evaluating the evidence base for the use of home telehealth remote monitoring in elderly with heart failure. *Telemed J E Health* 2009; **15**:783–796.
 56. Cleland JGF, Coletta AP, Clark AL. Clinical trials update from the European Society of Cardiology Heart Failure Meeting 2010: TRIDENT 1, BENEFICIAL, CUPID, RFA-HF, MUSIC, DUEL, Handheld BNP, phrenic nerve stimulation, CHAMPION and CABG with CRT study. *Eur J Heart Fail* 2010; **12**: 883–888.
 57. Jones NA, Frankel DS, Piette JD, Goldberg LR. Withdrawal of a technology-based daily weight monitoring system in patients with advanced heart failure eliminates mortality benefit (abstract). *Circulation* 2007; **116**:3277.